CORRECTION

Correction: Visceral adipose tissue mediates the relationship between left ventricular global longitudinal strain and insulin resistance among adults living with type 2 diabetes

Pavel Martinez-Dominguez^{1,2†}, Paola Gomez-Aviles^{3†}, Kenya Bautista-García^{4†}, Neftali Eduardo Antonio-Villa^{5†}, Enrique C. Guerra^{1,6}, Paloma Almeda-Valdes³, Alexandro J. Martagón^{3,7,8}, Alejandro Campos Munoz³, Maria Jose Santa-Ana-Bayona¹, Erick Alexanderson¹, Carlos A. Aquilar Salinas^{8,9} and Nilda Espinola-Zavaleta^{1,10*}

Correction to: Cardiovascular Diabetology (2025) 24:2ht tps://doi.org/10.1186/s12933-024-02547-x

In the original publication of this article [1], the open access funding statement was inadvertently included in Acknowledgements section. The following text "Open Access funding for this article was supported by National Institute of Cardiology Ignacio Chavez" has been removed from the article. The updated Acknowledgements section is shown below:

Acknowledgements

ECG is enrolled at the PECEM program of the Faculty of Medicine at UNAM and supported by CONACyT. The original article has been corrected.

[†]Pavel Martinez-Dominguez, Paola Gomez-Aviles, Kenya Bautista-García, Neftali Eduardo Antonio-Villa contributed equally to this work and were designated as co-first authors.

The original article can be found online at https://doi.org/10.1186/s1 2933-024-02547-x.

*Correspondence:

- Nilda Espinola-Zavaleta
- niesza2001@hotmail.com

BMC

- ¹Department of Nuclear Cardiology, National Institute of Cardiology Ignacio Chavez, Mexico City, Mexico
- ²Faculty of Medicine and Biomedical Sciences, Autonomous University of Chihuahua (UACH), Chihuahua, Mexico
- ³Endocrinology and Metabolism Department, Metabolic Research Unit, Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, Mexico

- ⁴School of Medicine, Benemérita Universidad Autónoma de Puebla, Puebla, Mexico
- ⁵Department of Endocrinology, Instituto Nacional de Cardiología Ignacio Chávez, Mexico City, Mexico
- ⁶Facultad de Medicina, Universidad Nacional Autónoma de México, Mexico City, Mexico
- ⁷The Institute for Obesity Research, Tecnologico de Monterrey, Mexico City, Mexico
- ⁸Escuela de Medicina y Ciencias de la Salud, Tecnologico de Monterrey, Mexico City, Mexico
- ⁹Instituto Nacional de Ciencias Médicas y Nutrición Salvador Zubirán, Mexico City, Mexico
- ¹⁰Echocardiography Department, ABC Medical Center, Mexico City, Mexico

© The Author(s) 2025. Open Access This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creati vecommons.org/licenses/by-nc-nd/4.0/.





Published online: 03 May 2025

Reference

Martinez-Dominguez P, Gomez-Aviles P, Bautista-García K, et al. Visceral adipose tissue mediates the relationship between left ventricular global longitudinal strain and insulin resistance among adults living with type 2 diabetes. Cardiovasc Diabetol. 2025;24:2.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.